Neuroablation for Intermetatarsal Nerve Entrapment (Morton's neuroma)

Clinical Policy ID: CCP.1370

Recent review date: 8/2026

Next review date: 12/2027

Policy contains: Achilles tendinosis; intermetatarsal nerve entrapment; Morton’s neuroma; plantar fasciitis; thermal ablation or chemoablation.

AmeriHealth Caritas has developed clinical policies to assist with making coverage determinations. AmeriHealth Caritas’s clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of “medically necessary,” and the specific facts of the particular situation are considered, on a case-by-case basis, by AmeriHealth Caritas when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. AmeriHealth Caritas’s clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients.’s AmeriHealth Caritas’s clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, AmeriHealth Caritaswill update its clinical policies as necessary.AmeriHealth Caritas’s clinical policies are not guarantees of payment.

Coverage policy

Neuroablation is clinically proven and, therefore, may be medically necessary to treat chronic foot pain caused by intermetatarsal nerve entrapment (also called Morton’s neuroma) (American Academy of Orthopaedic Surgeons, 2026).

Limitations

Not applicable

Alternative covered services

Standard of care for each diagnosis (e.g., analgesia medications, physical therapy, orthoses, surgery).

Background

Foot pain is a common complaint and includes diverse neurogenic and musculoskeletal causes, which often present with similar symptoms. The specific anatomic location of the pain (the forefoot, midfoot, and hindfoot) can often guide diagnosis and treatment. Morton's neuroma is a frequent cause of forefoot pain. It is a thickening of the interdigital nerve, commonly between the second/third and third/fourth toes. The exact cause is not known, but the inflammation is likely related to pressure on the nerve. Left untreated, nerve damage or chronic foot pain may occur (American Orthopaedic Foot and Ankle Society, undated).

The treatment algorithm for most chronically painful soft tissue conditions of the foot begins with conservative therapies (e.g., anti-inflammatory medications, off-loading, changes in footwear, orthoses, and self-care) and progresses to more invasive options. Prescribed options such as corticosteroid injections and physical therapy may relieve pain and improve mobility. When conservative treatment fails, surgery (e.g., decompression, release, or resection) may be indicated (Agyekum, 2015). Percutaneous techniques, such as platelet-rich plasma, extracorporeal shock wave therapy, and neuroablation have emerged as potential treatment options for chronic foot pain (Smith, 2017).

Neuroablation is designed to destroy neural tissue. In current practice, the goal of neuroablation is to temporarily interrupt rather than permanently destroy the nerve impulse or pathway, thus preventing the pain signal from traveling to the brain (Association of Extremity Nerve Surgeons, 2014). Neuroablation can be administered during open surgery or percutaneously with ultrasound, computed tomography, or magnetic resonance imaging guidance. The most common neuroablative techniques apply extreme heat (radiofrequency), cold (cryotherapy), or chemicals (e.g., ethanol) to denervate tissue.

The advantages of neuroablative techniques over surgery are minimal invasiveness, low associated morbidity, and repeatability should the pain recur. However, these methods damage tissue in a relatively blind manner without absolute control and may not permanently resolve symptoms (Association of Extremity Nerve Surgeons, 2014).

Findings

Guidelines

Evidence-based recommendations are conflicting on the value neuroablation especially for patients with intermetatarsal nerve entrapment (also called Morton’s neuroma), as professional societies attempt to balance the lack of quality evidence with increasing interest in nonsurgical, minimally invasive neuroablative approaches to treat chronic foot pain (American College of Foot and Ankle Surgeons [Schneider, 2018; Thomas, 2009]; Association of Extremity Nerve Surgeons, 2014).

For Morton’s neuroma, the American Academy of Orthopaedic Surgeons (2026) lists nerve-targeting procedures among nonsurgical treatment options for initial management, without preference. Their recommendations are based on studies showing that long-term relief can be achieved with a combination of shoe wear modifications, orthotics, anti-inflammatory medications (e.g., ibuprofen or naproxen), and/or corticosteroid injections. In some cases, minimally invasive nerve-targeting procedures may be used to reduce pain by disrupting nerve signaling:

- Alcohol (ethanol) injections chemically ablate the nerve to reduce pain signals.
- Radiofrequency ablation uses heat to create a controlled lesion in the nerve to reduce pain signals.
- Cryoablation uses cold to temporarily disrupt nerve function, although long-term outcomes are still being studied.

Evidence review

The evidence base consists of mostly retrospective case series of short-term duration (six months or less) and few comparative studies by which to compare the effectiveness of neuroablative techniques to surgical procedures considered the current standard of care.

The limitations in the evidence are those inherent to non-comparative, non-randomized, retrospective, and insufficiently powered studies. The literature provides low quality evidence supporting the safety and efficacy of radiofrequency ablation, cryoablation, and chemoablation (alcohol) for treating intermetatarsal nerve entrapment (Masala, 2018; Perini, 2016; Shah, 2019), plantar fasciitis (Adán-Laguna, 2026; Kurtoglu, 2022; Landsman, 2013), and Achilles tendinosis (Wei, 2017). These procedures appear safe and efficacious in the short-term with durable results in some instances, but the optimal patient selection criteria or ablative protocol cannot be determined.

Systematic reviews expose the lack of high-quality evidence comparing the effectiveness of available treatment options, including neurolytic methods, for Morton's neuroma (Lu, 2021; Matthews, 2019; Santos, 2018; Thomson, 2020). More recently, Llombart-Blanco's 2025 systematic review of eight studies (n = 237) found radiofrequency ablation achieved significant pain relief in 47.57% of participants (95% confidence interval 25.13% to 70.00%), particularly at temperatures ≥ 85°C and with fewer radiofrequency cycles (≤ three). Complications were minimal (2.1%), and most resolved without significant interventions.

An incremental cost-utility analysis concluded that the most cost-effective approach to managing Morton's neuroma after conservative therapy has failed is beginning with the least expensive option (steroid injection) and proceeding only when necessary in a stepwise manner to more invasive and expensive treatments, such as ultrasound-guided alcohol ablation, and then surgery (Ross, 2022).

In 2019, we updated the references. No policy changes are necessary. The policy ID was changed from CP# 14.02.14 to CCP.1317.

In 2020, we updated the references. No policy changes are warranted.

In 2021, we updated the references. No policy changes are warranted.

In 2022, we updated the references with no policy changes warranted.

In 2023, we updated the references with no policy changes warranted.

In 2024, no new relevant publications were found and no policy changes were warranted.

In 2025, we updated the references with no policy changes warranted.

In 2026, we updated the references, modified the title to address intermetatarsal nerve entrapment specifically, and modified medical necessity criteria to align with American Academy of Orthopaedic Surgeons latest recommendations.

References

On August 18, 2026, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were "Ablation technique" (MeSH), "Metatarsalgia" (MeSH), "Peripheral nerves" (MeSH), "neuroablation," "radiofrequency ablation," "cryoablation," "plantar," and "foot." We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

Adán-Laguna J, Villarroya Martínez L, Puyuelo Martínez G, et al. Radiofrequency neurolysis versus surgical neurectomy for Morton's neuroma. *Int Orthop*. 2026. Doi: 10.1007/s00264-026-06824-0.

Agyekum EK, Ma K. Heel pain: A systematic review. *Chin J Traumatol.* 2015;18(3):164-169. Doi: 10.1016/j.cjtee.2015.03.002.

American Academy of Orthopaedic Surgeons. OrthoInfo. Diseases & Conditions. Morton's neuroma. https://orthoinfo.aaos.org/en/diseases--conditions/mortons-neuroma. Published 2026.

American Orthopaedic Foot and Ankle Society. FootCareMD. Conditions and treatments. https://www.footcaremd.org/foot-and-ankle-conditions/toes/mortons-neuroma . Undated.

Association of Extremity Nerve Surgeons. Clinical practice guidelines. Edition 1. Denervation. https://www.aens.us/images/aens/AENS%20Guidelines%20-%20Final.pdf. Published November 2014.

Kurtoglu A, Kochai A, Inanmaz ME, et al. Effectiveness of radiofrequency ablation for treatment of plantar fasciitis. *Medicine (Baltimore)*. 2022;101(12):e29142. Doi: 10.1097/MD.0000000000029142.

Landsman AS, Catanese DJ, Wiener SN, Richie DH, Jr., Hanft JR. A prospective, randomized, double-blinded study with crossover to determine the efficacy of radio-frequency nerve ablation for the treatment of heel pain. *J Am Podiatr Med Assoc.* 2013;103(1):8-15. Doi: 10.7547/10300008.

Llombart-Blanco R, Mariscal G, Benlloch M, Barrios C, Llombart-Ais R. Systematic review and meta-analysis of radiofrequency ablation for Morton's neuroma: Outcomes and predictors of success. *Am J Phys Med Rehabil*. 2025;104(5):465-472. Doi: 10.1097/phm.0000000000002668.

Lu VM, Puffer RC, Everson MC, et al. Treating Morton's neuroma by injection, neurolysis, or neurectomy: A systematic review and meta-analysis of pain and satisfaction outcomes. *Acta Neurochir (Wien).* 2021;163(2):531-543. Doi: 10.1007/s00701-020-04241-9.

Masala S, Cuzzolino A, Morini M, Raguso M, Fiori R. Ultrasound-guided percutaneous radiofrequency for the treatment of Morton's neuroma. *Cardiovasc Intervent Radiol*. 2018;41(1):137-144. Doi: 10.1007/s00270-017-1786-y.

Matthews BG, Hurn SE, Harding MP, Henry RA, Ware RS. The effectiveness of non-surgical interventions for common plantar digital compressive neuropathy (Morton's neuroma): A systematic review and meta-analysis. *J Foot Ankle Res.* 2019;12:12. Doi: 10.1186/s13047-019-0320-7.

Perini L, Perini C, Tagliapietra M, et al. Percutaneous alcohol injection under sonographic guidance in Morton's neuroma: Follow-up in 220 treated lesions. *Radiol Med.* 2016;121(7):597-604. Doi: 10.1007/s11547-016-0622-9.

Ross AB, Jacobs A, Williams KL, Bour RK, Gyftopoulos S. Ultrasound-guided injection treatments versus surgical neurectomy for Morton Neuroma: A cost-effectiveness analysis. *AJR Am J Roentgenol*. 2022;218(2):234-240. Doi: 10.2214/AJR.21.26419.

Santos D, Morrison G, Coda A. Sclerosing alcohol injections for the management of intermetatarsal neuromas: A systematic review. *Foot (Edinb).* 2018;35:36-47. Doi: 10.1016/j.foot.2017.12.003.

Schneider HP, Baca JM, Carpenter BB, Dayton PD, Fleischer AE, Sachs BD. American College of Foot and Ankle Surgeons clinical consensus statement: Diagnosis and treatment of adult acquired infracalcaneal heel pain. *J Foot Ankle Surg*. 2018;57(2):370-381. Doi: 10.1053/j.jfas.2017.10.018.

Shah R, Ahmad M, Hanu-Cernat D, Choudhary S. Ultrasound-guided radiofrequency ablation for treatment of Morton's neuroma: Initial experience. *Clin Radiol*. 2019;74(10):815.e9-815.e13. Doi: 10.1016/j.crad.2019.07.002.

Smith WB, Melton W, Davies J. Midsubstance tendinopathy, percutaneous techniques (platelet-rich plasma, extracorporeal shock wave therapy, prolotherapy, radiofrequency ablation). *Clin Podiatr Med Surg.* 2017;34(2):161-174. Doi: 10.1016/j.cpm.2016.10.005.

Thomas JL, Blitch EL4th, Chaney DM, et al. Diagnosis and treatment of forefoot disorders. Section 3. Morton's intermetatarsal neuroma. *J Foot Ankle Surg*. 2009;48(2):251-256. Doi: 10.1053/j.jfas.2008.12.005.

Thomson L, Aujla RS, Divall P, Bhatia M. Non-surgical treatments for Morton's neuroma: A systematic review. *Foot Ankle Surg.* 2020;26(7):736-743. Doi: 10.1016/j.fas.2019.09.009.

Wei M, Liu Y, Li Z, Wang Z. Comparison of clinical efficacy among endoscopy-assisted radio-frequency ablation, extracorporeal shockwaves, and eccentric exercises in treatment of insertional Achilles tendinosis. *J Am Podiatr Med Assoc.* 2017;107(1):11-16. Doi: 10.7547/14-146.

Policy updates

3/2018: initial review date and clinical policy effective date: 5/2018

5/2019: Policy references updated. Policy ID changed.

6/2020: Policy references updated.

6/2021: Policy references updated.

6/2022: Policy references updated.

6/2023: Policy references updated.

6/2024: Policy references updated.

6/2025: Policy references updated.

6/2026: Policy references updated.

9/2026: Policy references updated. Coverage modified.

Related codes

Below are the most commonly submitted codes for the service(s)/item(s) subject to this policy CCP.1370. This is not an exhaustive list of codes. Providers are expected to consult the appropriate coding manuals and bill accordingly.

Code	Code description
64632	Destruction by neurolytic agent; plantar common digital nerve
0441T	Ablation, percutaneous, cryoablation, includes imaging guidance; lower extremity distal/peripheral nerve