

# Erectile dysfunction treatments other than pharmaceuticals

Clinical Policy ID: CCP.5555

Policy Contains: surgical revascularization; penile vacuum pump; erectile dysfunction; penile prosthesis; impotence

Recent review date: 12/3/2019

Next review date: 4/2021

---

## Coverage policy

---

Use of vacuum erection devices, penile prosthesis implantation, and penile arterial reconstruction for erectile dysfunction are clinically proven and, therefore, medically necessary when the following criteria are met:

- The member is diagnosed with erectile dysfunction.
- Conservative treatments have been attempted for at least 12 months and have failed (Burnett, 2018).

Removal of a penile implantation is considered medically necessary for an infected prosthesis, intractable pain, mechanical failure, or urinary obstruction.

Reimplantation of a penile implant is considered medically necessary for members whose prior prosthesis was removed for medically necessary indications.

### Limitations

All other nonmedicinal interventions for erectile dysfunction are considered not clinically proven, and therefore, not medically necessary — including venous surgery, low-intensity extracorporeal shock wave therapy, intra-cavernosal stem cell therapy, and platelet-rich plasma therapy (Burnett, 2018).

### Alternative covered services

Various medications (not addressed in this policy).

---

## Background

---

Erectile dysfunction, also referred to as impotence, is defined as the inability to achieve or maintain an erection that is sufficient for satisfactory sexual performance. The condition is more common in older males — one random sample placed the prevalence of minimal, moderate, and complete impotence for males ages 40 – 70 at 52% (O'Donnell, 2004). The crude annual incidence rate of the condition is estimated at 25.9 per 1,000 man-years, with incidence rising with each decade of life (Johannes, 2000).

Erectile dysfunction was once believed to be a psychological disease, but more than 80% of cases are now considered to have an organic etiology. Conditions associated with the disorder include hypogonadism, lower urinary tract symptoms, benign prostatic hypertrophy, hypertension, cardiovascular disorder, smoking, excess alcohol intake, obesity, dyslipidemia, diabetes mellitus, metabolic syndrome, stress, anxiety, and depression. Reactions to various surgeries and medications can also cause erectile dysfunction (Yafi, 2016).

Diagnosing erectile dysfunction includes a work-up of patients seeking medical care for the disorder. The diagnosis can only be made with establishment of an accurate medical and sexual history; a careful general and focused genitourinary examination; and a minimum number of hormonal and routine biochemical tests. Provider questioning to the patient must be done in a manner to minimize patient embarrassment (Yafi, 2016).

Erectile dysfunction is a highly under-treated condition. A study of 6.2 million males diagnosed with erectile dysfunction found that only 25.4% were treated (at least one filled prescription for phosphodiesterase type 5 inhibitor, injection or urethral prostaglandins, or androgen replacement) over a 12-month period. Men older than age 60 were significantly less likely ( $P < .0001$ ) to be treated than males ages 40 – 59 (Frederick, 2014).

Recommended treatment of men with erectile dysfunction should always include encouragement of patient lifestyle changes that address known causes of the disorder. These include changes in diet, increased physical activity, and cessation of alcohol consumption or tobacco use. The most conservative first-line treatment for the disorder is prescribed oral phosphodiesterase type 5 inhibitors; testosterone therapy can be added if the patient also has hypogonadism (Burnett, 2018).

When conservative treatments do not result in improvements, erectile dysfunction can be treated using several more invasive approaches, namely:

- Penile Prosthesis. Inflatable pumps or semi-rigid/malleable rods are types of prostheses that can be implanted into the penis and scrotum in a one- to two-hour procedure. Infections can occur, along with complications such as glans bowing, reservoir complications, corporal crossover, and perforations (Sadeghi-Nejad, 2013).
- Vacuum Erection Device. An acrylic cylinder with a pump may be attached directly to the end of the penis, and a constriction ring or band is placed on the cylinder at the other end, which is applied to the body. The cylinder and pump create a vacuum to help the penis become erect, while the band or constriction ring helps maintain the erection (Hoyland, 2013).
- Penile Arterial Reconstruction. Vascular surgery can reconstruct arteries to improve blood flow to the penis. Recent types of this surgery include circumferential incision plus a median pubic longitudinal approach with acupuncture-assisted local anesthesia and penile venous stripping surgery (Molodysky, 2013).

Dietary supplements and other natural treatments are also used to treat erectile dysfunction; efficacy reviews in the professional literature are limited.

---

## Findings

---

The American Urological Association guideline on erectile dysfunction recommended six types of treatments that could be considered. Three are medications, while the others are vacuum erection devices, penile prosthesis implantation, and penile arterial reconstruction — for each, the guideline states that patients must

be informed of potential risks and benefits before treatment starts. Treatments not recommended are venous surgery, low-intensity extracorporeal shock wave therapy, intracavernosal stem cell therapy, and platelet-rich plasma therapy (Burnett, 2018).

The Canadian Urological Association agreed that oral medications should be first-line therapy. However, second-line therapies and surgery are also important options in treating confirmed cases of erectile dysfunction (Bella, 2015). The European Association of Urology also supported these first-, second-, and third-line therapies (Hatzimouratidis, 2016).

The American Academy of Family Physicians guideline recommended lifestyle changes (including tobacco cessation, exercise, weight loss, control of diabetes, hypertension, and hyperlipidemia), plus oral phosphodiesterase-5 inhibitors as first-line treatments for erectile dysfunction. The Academy recommended alprostadil and vacuum devices for second-line therapy, and surgically implanted penile prostheses when other treatments have failed (Rew, 2016).

The American Society of Clinical Oncology endorsed a guideline, including a recommendation that people with cancer be counseled about sexual health and dysfunction related to cancer. The guideline states that if medical management does not succeed, medication such as phosphodiesterase type 5 inhibitors may be beneficial, and surgery remains an option for males with erectile dysfunction (Carter, 2018).

A number of systematic reviews or meta-analyses and other large-scale studies have appeared in the professional literature addressing safety and effectiveness of various treatments for erectile dysfunction.

### Penile Prosthesis

- A systematic review of 19 articles consisting of males undergoing penile prosthesis placement found organic conditions caused almost all cases of erectile dysfunction. Average duration of the condition before the procedure was 56 months and 72 months for those undergoing inflatable and malleable prostheses, respectively. People with diabetes undergoing the inflatable procedure had an average of 75 months (Bajic, 2019).
- A systematic review of 14 studies ( $n = 9,910$  patients with a first-time penile implant) determined that the infection rate for patients whose prosthesis did not have an infection retardant coating was significantly greater than those who had such a coating (2.32% versus 0.89%, or  $P < .01$ ) (Mandava, 2012).
- An extensive review of the professional literature identified several advanced and novel techniques for penile lengthening that have been successfully performed at the time of insertion of a penile prosthesis in males with erectile dysfunction (Tran, 2017).
- A retrospective study of 883 patients with erectile dysfunction who underwent malleable, two-piece, or three-piece penile prosthesis implants and were followed for an average of 49 months, showed couples' satisfaction was significantly highest in the two- and three-piece group. The highest rate of revision surgery due to penile corporal perforation was in the malleable group ( $P = .021$ ), whereas the highest rate of revision surgery due to penile implant malfunction occurred in the three-piece implant group ( $P = .001$ ) (Cayan, 2019).
- In a review of only included studies with at least a five-year follow up, five- and 10-year device survival of prosthesis implant was 90.4% and 86.6%, respectively. Eight- and 10-year infection rates were 1.5% and 1.8%. Authors maintain that prosthesis implant is the gold standard for erectile dysfunction patients refractory to medicinal treatment (Dick, 2019).

## Vacuum Erection Device

- A systematic review or meta-analysis of six controlled trials ( $n = 273$ ) assessed vacuum therapy for penile rehab after radical prostatectomy. Early use of vacuum therapy significantly improved erectile function and penile shrinkage. Few adverse events and no serious side effects were reported (Qin, 2018).
- A study of 141 sexually active patients who underwent radical prostatectomy with early use of a vacuum erection device as a prophylaxis showed that after five years, 62% remained sexually active, of whom 71% had natural erections not requiring assistance (Raina, 2010).
- A survey answered by 618 urologists on treatments for erectile dysfunction following radical prostatectomy showed the vacuum erection device was the most-commonly used (nondrug) primary strategy (Tal, 2011).
- A study of 1,500 men with organic erectile dysfunction participated in vacuum constrictive device training. At the first session, 87.4% attained a full erection, while the others did so after one week. The ability to perform vaginal penetration was 94.6%. Erectile scores of improved from 9.3 to 27.5 ( $P < .05$ ) (Khayyamfar, 2014).

## Penile Arterial Reconstruction

- A review of 25 nonrandomized studies of penile revascularization surgery found the subjective cure rate in men under 30 was better than older men ( $P = .001$ ). Venous leak and history of smoking influenced success rates (Babaei, 2009).
- A systematic review or meta-analysis of 16 articles ( $n = 374$ ) assessed efficacy and safety of endovascular therapy in patients with veno-occlusive dysfunction or arterial insufficiency. Overall clinical success rates for the groups were 59.8% and 63.2%; complications occurred in 5.2% and 4.9% (Doppalapudi, 2019).
- A study of 96 patients with (veno-occlusive) erectile dysfunction found pelvic venoablation, without any drugs, allowed 80.21% to have erections sufficient for vaginal insertion within three months (Herwig, 2015).
- A study of 110 patients tracked an average of 73.2 months after penile revascularization surgery showed an increase in erection function from 7.3 to 16.8. The three-month success rate ( $> 5$ -point increase), was 81.8% at three months, and 63.6% at five years (Kayigil, 2012).

## Extracorporeal shock wave therapy

- A meta-analysis of seven controlled trials ( $n = 522$ ) compared low-intensity extracorporeal shock wave therapy for erectile dysfunction with sham therapy. Significant improvements for the treatment group were observed for erectile function ( $P < .00001$ ), while the best improvements were for moderate or severe erectile dysfunction. Follow up in five of seven studies were five weeks or less; in the other two, which followed subjects for 12 months, only one showed greater efficacy after treatment (Dong, 2019).
- A systematic review of low-intensity shock wave therapy for erectile dysfunction included 11 studies ( $n = 799$ ). Nine studies found a significant improvement in erectile function after six months. However, after 12 months, two of five studies found a plateauing, and the other three a deterioration (Brunckhourst, 2019).
- A meta-analysis of 10 controlled trials ( $n = 873$ ) found low-intensity extracorporeal shock wave therapy for erectile dysfunction improved function ( $P = .0009$ ) and patient outcomes ( $P < .00001$ ) (Sokolakis, 2019).

- A systematic review or meta-analysis documented that, based on nine studies (n = 637), low-energy extracorporeal shock wave therapy significantly improved erection function ( $P = .003$ ) for three months (Man, 2018).
- A systematic review or meta-analysis of 15 studies (n = 277) found low-intensity extracorporeal shock wave treatment was 8.31 times more effective than sham treatment, and improved erectile function by 2.5 times more, one month after treatment (Zou, 2017).
- A systematic review or meta-analysis of 14 trials (n = 833) showed low-intensity extracorporeal shock wave therapy improved erection function ( $P < .0001$ ) and erection hardness score ( $P < .01$ ), and that efficacy could last at least three months (Lu, 2017).

#### Other treatments

- A review of 19 articles addressing relatively new interventions for erectile dysfunction documented evidence supporting the use of two microsurgical treatments, namely microvascular arterial bypass penile revascularization surgery and cavernous nerve graft reconstruction (Shaully, 2019).

A review of the literature on erectile function following radical prostatectomy could not arrive on one type of treatment that was superior to others (Mulhall, 2013).

---

## Billing and coding

---

Below are National Coverage Determinations, Local Coverage Determinations, and the most commonly submitted codes subject to this policy. This is not an exhaustive list of codes. Providers are expected to consult the appropriate Centers for Medicare & Medicaid Services references and coding manuals, and bill accordingly.

#### National coverage determinations

No National Coverage Determinations were identified as of the writing of this policy.

#### Local coverage determinations

No Local Coverage Determinations were identified as of the writing of this policy.

#### CPT procedure codes

[ccp.28B\_CPT]

#### HCPCS level II

[ccp.28C\_HCPCSII]

#### ICD-10 diagnostic codes

[ccp.28A\_ICD10]

---

## References

---

On August 15, 2019, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “erectile dysfunction,” “extracorporeal shock wave therapy,” “penile arterial reconstruction,” “penile prosthesis implantation,” and “vacuum erection devices.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

- Babaei AR, Safarinejad MR, Kolahi AA. Penile revascularization for erectile dysfunction: a systematic review and meta-analysis of effectiveness and complications. *Urol J*. 2009;6(1):1-7. <https://www.ncbi.nlm.nih.gov/pubmed/?term=%22erectile+dysfunction%22+venous+surgery>. Accessed August 19, 2019.
- Bajic P, Mahon J, Faraday M, Sadeghi-Nejad H, Hakim L, McVary KT. Etiology of erectile dysfunction and duration of symptoms in patients undergoing penile prosthesis: A systematic review. *Sex Med Rev*. 2019 Jul 2. pii: S2050-0521(19)30042-3. Doi: 10.1016/j.sxmr.2019.05.003.
- Bella AJ, Lee JC, Carrier S, Benard F, Brock GB. 2015 CUA practice guidelines for erectile dysfunction. *Can Urol Assoc J*. 2015;9(1-2):23–29. Doi: 10.5489/cuaj.2699.
- Brunckhorst O, Wells L, Teeling F, Muir G, Muneer A, Ahmed K. A systematic review of the long-term efficacy of low-intensity shock wave therapy for vasculogenic erectile dysfunction. *Int Urol Nephrol*. 2019;51(5):773-781. Doi: 10.1007/s11255-019-02127-z
- Burnett AL, Nehra A, Breau RH, et al. Erectile dysfunction: AUA guideline (2018). American Urological Association. [https://www.auanet.org/guidelines/erectile-dysfunction-\(ed\)-guideline](https://www.auanet.org/guidelines/erectile-dysfunction-(ed)-guideline). Published 2018. Accessed August 15, 2019.
- Carter J, Lacchetti C, Andersen BL, et al. Interventions to address sexual problems in people with cancer: American Society of Clinical Oncology clinical practice guideline adaptation of Cancer Care Ontario guideline. *J Clin Oncol*. 2018;36(5):492-511. Doi: 10.1200/JCO.2017.75.8995.
- Cayan S, Asci R, Efesoy O, Bolat MS, Akbay E, Yaman O. Comparison of long-term results and couples' satisfaction with penile implant types and brands: Lessons learned from 883 patients with erectile dysfunction who underwent penile prosthesis implantation. *J Sex Med*. 2019;16(7):1092-1099. Doi: 10.1016/j.jsxm.2019.04.013.
- Dick B, Tsambarlis P, Reddy A, Hellstrom WJ. An update on: long-term outcomes of penile prostheses for the treatment of erectile dysfunction. *Expert Rev Med Devices*. 2019;16(4):281-286. Doi: 10.1080/17434440.2019.1598259.
- Dong L, Chang D, Zhang X, et al. Effect of low-intensity extracorporeal shock wave on the treatment of erectile dysfunction: A systematic review and meta-analysis. *Am J Mens Health*. 2019;13(2):1557988319846749. Doi: 10.1177/1557988319846749.
- Doppalapudi SK, Wajswol E, Shukla PA, et al. Endovascular therapy for vasculogenic erectile dysfunction: A systematic review and meta-analysis of arterial and venous therapies. *J Vasc Interv Radiol*. 2019;30(8):1251-1258.e2. Doi: 10.1016/j.jvir.2019.01.024.
- Frederick LR, Cakir OO, Arora H, Halfand BT, McVary KT. Undertreatment of erectile dysfunction: claims analysis of 6.2 million patients. *J Sex Med*. 2014;11(10):2546-2553. Doi: 10.1111/jsm.12647.
- Hatzimoutidis K, Giuliano F, Moncada I, Muneer A, Salonia A, Verge P. EAU guidelines on erectile dysfunction, premature ejaculation, penile curvature, and priapism. European Association of Urology. <https://uroweb.org/wp-content/uploads/EAU-Guidelines-Male-Sexual-Dysfunction-2016-3.pdf>. Published 2016. Accessed August 15, 2019.

Herwig R, Sansalone S. Venous leakage treatment revisited: pelvic venoablation using aethoxysclerol under air block technique and Valsalva maneuver. *Arch Ital Urol Androl*. 2015;87(1):1-4. Doi: 10.4081/aiua.2015.1.1.

Hoyland K, Vasdev N, Adshead J. The use of vacuum erection devices in erectile dysfunction after radical prostatectomy. *Rev Urol*. 2013;15(2):67-71. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3784970/>. Accessed August 19, 2019.

Johannes CB, Araujo AB, Feldman HA, Derby CA, Kleinman KP, McKinlay JB. Incidence of erectile dysfunction in men 40 to 69 years old: longitudinal results from the Massachusetts male aging study. *J Urol*. 2000;163(2):460-463. <https://www.ncbi.nlm.nih.gov/pubmed/10647654>. Accessed August 16, 2019.

Kayigil O, Okulu E, Aldemir M, Onen E. Penile revascularization in vasculogenic erectile dysfunction (ED): long-term follow-up. *BJU Int*. 2012;109(1):109-115. Doi: 10.1111/j.1464-410X.2011.10293.x.

Khayyamfar F, Forootan SK, Ghasemi H, Miri SR, Farhadi E. Evaluating the efficacy of vacuum constrictive device and causes of its failure in impotent patients. *Urol J*. 2014;10(4):1072-1078. <https://www.ncbi.nlm.nih.gov/pubmed/?term=Kheyamfar+F%2C+Forootan+SK%2C+Ghasemi+H>. Accessed August 21, 2019.

Li P, Shen YJ, Liu TQ, et al. Vacuum therapy for erectile dysfunction that fails to respond to PDE-5i: report of 70 cases. *Zhonghua Nan Ke Xue*. 2013;19(3):236-240. <https://www.ncbi.nlm.nih.gov/pubmed/23700730>. Accessed August 20, 2019.

Lu Z, Lin G, Reed-Maldonado A, Wang C, Lee YC, Lue TF. Low-intensity extracorporeal shock wave treatment improves erectile function: A systematic review and meta-analysis. *Eur Urol*. 2017;71(2):223-233. Doi: 10.1016/j.eururo.2016.05.050.

Man L, Li G. Low-intensity extracorporeal shock wave therapy for erectile dysfunction: A systematic review and meta-analysis. *Urology*. 2018;119:97-103. Doi: 10.1016/j.urology.2017.09.011.

Mandava SH, Serefoglu EC, Freier MT, Wilson SK, Hellstrom WJ. Infection retardant coated inflatable penile prostheses decrease the incidence of infection: a systematic review and meta-analysis. *J Urol*. 2012;188(5):1855-60. Doi: 10.1016/j.juro.2012.07.022.

Molodysky E, Liu S-P, Huang S-J, Hsu G-L. Penile vascular surgery for treating erectile dysfunction: Current role and future direction. *Arab J Urol*. 2013;11(3):254-266. Doi: 10.1016/j.aju.2013.05.001.

Mulhall JP, Bivalacqua TJ, Becher EF. Standard operating procedure for the preservation of erectile function outcomes after radical prostatectomy. *J Sex Med*. 2013;10(1):195-203. Doi: 10.1111/j.1743-6109.2012.02885.x.

O'Donnell AB, Araujo AB, McKinlay JB. The health of normally aging men: The Massachusetts Male Aging Study (1987-2004). *Exp Gerontol*. 2004;39(7):975-984. H. <https://www.ncbi.nlm.nih.gov/pubmed/15236757>. Accessed August 21, 2019.

Qin F, Wang S, Li J, Wu C, Yuan J. The early use of vacuum therapy for penile rehabilitation after radical prostatectomy: Systematic review and meta-analysis. *Am J Mens Health*. 2018;12(6):2136-2143. Doi: 10.1177/1557988318797409.

Raina R, Pahlajani G, Agarwal A, Jones S, Zippe C. Long-term potency after early use of a vacuum erection device following radical prostatectomy. *BJU Int*. 2010;106(11):1719-1722. Doi: 10.1111/j.1464-410X.2010.09360.x.

Rew KT, Heidelbaugh JJ. Erectile dysfunction. *Am Fam Physician*. 2016;94(10):820-827. <https://www.ncbi.nlm.nih.gov/pubmed/27929275>. Accessed August 15, 2019.

Sadeghi-Nejad H, Fam M. Penile prosthesis surgery in the management of erectile dysfunction. *Arab J Urol*. 2013;11(3):245-253. Doi: 10.1016/j.aju.2013.05.002.

Shaully O, Gould DJ, Patel KM. Emerging nonsurgical and surgical techniques to treat erectile dysfunction: A systematic review of treatment options and published outcomes. *J Plast Reconstr Aesthet Surg*. 2019;72(4):532-538. Doi: 10.1016/j.bjps.2018.12.028.

Sokolakis I, Hatzichristodoulou G. Clinical studies on low intensity extracorporeal shock wave therapy for erectile dysfunction: a systematic review and meta-analysis of randomized controlled trials. *Int J Impot Res*. 2019;31(3):177-194. Doi: 10.1038/s41443-019-0117-z.

Tal R, Teloken P, Mulhall JP. Erectile function rehabilitation after radical prostatectomy: practice patterns among AUA members. *J Sex Med*. 2011;8(8):2370-2376. Doi: 10.1111/j.1743-6109.2011.02355.x.

Tran H, Goldfarb R, Ackerman A, Valenzuela RJ. Penile lengthening, girth, and size preservation at the time of penile prosthesis insertion. *Sex Med Rev*. 2017;5(3):403-412. Doi: 10.1016/j.sxmr.2016.11.005.

Yafi FA, Jenkins L, Albersen M, et al. Erectile dysfunction. *Nat Rev Dis Primers*. 2016;2:16003. Doi: 10.1038/nrdp.2016.3.

Yang G, Muzepper M. Platelet indices and erectile dysfunction: A systematic review and meta-analysis. *Andrologia*. 2019;51(5):e13248. Doi: 10.1111/and.13248.

Zou ZJ, Tang LY, Liu ZH, et al. Short-term efficacy and safety of low-intensity extracorporeal shock wave therapy in erectile dysfunction: A systematic review and meta-analysis. *Int Braz J Urol*. 2017;43(5):805-821. Doi: 10.1590/S1677-5538.IBJU.2016.0245.

---

## Policy updates

---

Initial review date: 12/3/2019

Effective date: 1/1/2020

No policy updates.